

#### CHARACTERISTICS

RATTO MORTO BLOCK is a ready for use wax block rodenticide bait containing an anticoagulant active substance, Brodifacoum, effective against House mouse (*Mus musculus*), Brown rat (*Rattus norvegicus*), after single ingestion. Formulation consist of a wax block appetizing for target organisms. The product does not alert or generate suspicions in rodent population. RATTO MORTO BLOCK contains a bittering agent to prevent accidental ingestion from children. RATTO MORTO BLOCK can be used indoor or outdoor, around building as home, cellars, garages, storage rooms and private gardens.

#### TARGET ORGANISMS

House mouse (*Mus musculus*), juveniles to adults.  
Brown rat (*Rattus norvegicus*), juveniles to adults.

#### APPLICATION METHOD

RATTO MORTO BLOCK is a ready-to-use wax block bait to be used in tamper-resistant bait stations, labelled, protected from atmospheric conditions, ingestion of non-target species and dispersion in the environment.

#### INSTRUCTIONS FOR USE

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it. Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered. Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve. Bait stations should be placed in the immediate vicinity where rodent activity has been observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.). Where possible, bait stations must be fixed to the ground or other structures. Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals. Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these. Do not place bait stations near water drainage systems where they can come into contact with water. When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product. Remove the remaining bait or the bait stations at the end of the treatment period.

#### RISK MITIGATION MEASURES

Do not handle until all safety precautions have been read and understood. Do not eat, drink or smoke when using this product. Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinfestation. Do not use anticoagulant rodenticides as permanent baits (e.g. for prevention of rodent infestation or to detect rodent activity).

The product information (i.e. label and/or leaflet) shall clearly show that:  
- the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").  
- users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. "label bait stations according to the product recommendations").

Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service. Search for and remove dead rodents during treatment, at least as often as bait stations are inspected. Dispose dead rodents in accordance with local requirements.

Poisoned rodents should be double-bagged using plastic bags and either disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets or collected by a specialist waste contractor or the local authority.

#### PARTICULARLY OF LIKELY DIRECT OR INDIRECT EFFECTS, FIRST AID INSTRUCTIONS AND EMERGENCY MEASURES TO PROTECT ENVIRONMENT

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

Antidote: Vitamin K1 administered by medical/veterinary personnel only.

In case of:

- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet.

UK medical professionals should contact the National Poisons Information Service ([www.npis.org](http://www.npis.org)) for further advice.  
Hazardous to wildlife.

#### INSTRUCTION FOR SAFE DISPOSAL OF THE PRODUCT AND ITS PACKAGING


At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements. In the UK waste bait should be double bagged in plastic bags and disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets or taken to a civic amenity site. For information on civic amenity sites contact the local authority.

#### CONDITIONS OF STORAGE AND SHELF-LIFE OF THE PRODUCT UNDER NORMAL CONDITIONS OF STORAGE

Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight. Store in places prevented from the access of children, birds, pets and farm animals.

Shelf life: 2 years.

# RATTO MORTO BLOCK



Net Content

260 g

deeval

Deeval International

HSE Authorisation no: UK-2019-1227

#### DOSES AND METHOD OF USE

##### Indoor against House mouse (*Mus musculus*).

Place 20 - 40 g of RATTO MORTO BLOCK per bait station, 1 - 2 unit of 20 g bait or 2 - 4 unit of 10 g bait. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

##### Indoor against Brown rat (*Rattus norvegicus*).

Place 100 - 200 g of RATTO MORTO BLOCK per bait station, 5 - 10 unit of 20 g bait or 10 - 20 unit of 10 g bait. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. The bait stations should be visited at least every 5 to 7 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

##### Outdoor, around buildings against Brown rat (*Rattus norvegicus*).

Place 100 - 200 g of RATTO MORTO BLOCK per bait station, 5 - 10 unit of 20 g bait or 10 - 20 unit of 10 g bait. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. Place the bait stations in areas not liable to flooding. Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt. The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

#### OTHER INFORMATION

Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait. Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them. This product contains a bittering agent and a dye.

#### RATTO MORTO BLOCK

Ready-to-use wax block rodenticide bait  
Category of users: **GENERAL PUBLIC**  
Field of use: **INDOOR and OUTDOOR, AROUND BUILDINGS**

#### Composition:

Brodifacoum	0,0025 %
(CAS N° 56073-10-0)	
Denatonium benzoate	0,001 %
(CAS N° 3734-33-6)	
Non-active substances	up to 100 %

#### WARNING



#### HAZARD STATEMENTS

H373 May cause damage to organs (blood) through prolonged or repeated exposure.

#### PRECAUTIONARY STATEMENTS

P102 Keep out of reach of children.  
P301 + P310 IF SWALLOWED: Immediately call a POISON CENTER or a doctor/physician/. P308 + P313 IF exposed or concerned: Get medical advice. P501 Dispose of contents to hazardous waste facilities in accordance with national regulations.

#### BIOCIDAL PRODUCT (PT14)

HSE Authorisation no: UK-2019-1227

#### Authorization holder:

DEEVAL INTERNATIONAL LTD  
Unit 45, First Avenue - Deeside Industrial Park CH5 2NU Flintshire Unite Kingdom - Tel. +44 07798714948

Packaging: 260g, containing wax block bait in 20g unit.

Shelf-life: 2 years

Batch n. / prod. date: